

Bevacizumab in Combination with Chemotherapy or Molecularly Targeted Agents for Non-Small-Cell Lung Cancer with Brain Metastases

To the Editor:

Su and Rau¹ have recently reported a case of non-small-cell lung cancer (NSCLC) of great interest. The patient, with anaplastic lymphoma kinase (ALK)-positive adenocarcinoma of the lung with brain metastases, was initially treated with whole-brain irradiation and then treated with chemotherapy consisting of pemetrexed and cisplatin; however, her brain metastases increased in size accompanied by muscle weakness and myoclonic jerks after two cycles of chemotherapy. The

authors then added bevacizumab to the same chemotherapy and the brain metastases markedly shrank.

To date, it has been prospectively demonstrated that bevacizumab is safe and effective in NSCLC patients with brain metastases as long as these metastases are pretreated² or asymptomatic.³ Moreover, a case series has shown that bevacizumab is also safe and effective for NSCLC patients with symptomatic, heavily pretreated brain metastases.⁴ We also experienced a case of lung adenocarcinoma in which the brain metastases were refractory to whole-brain irradiation but dramatically responded to combination chemotherapy of bevacizumab, paclitaxel, and carboplatin (Fig. 1). These findings suggest that bevacizumab may merit further investigation in NSCLC patients with symptomatic, heavily pretreated, and/or radioresistant brain metastases.

In addition, a recent retrospective study demonstrated that brain metastases are considerably common at the time of diagnosis and also at the time of disease progression in epidermal growth factor receptor (EGFR)-positive or ALK-positive NSCLC treated with tyrosine kinase inhibitors (TKIs).⁵ Considering that bevacizumab may have some booster effect for brain metastases and that brain metastases have a negative impact on survival in EGFR-positive NSCLC treated with EGFR-TKI as well,⁶ prospective

evaluation of bevacizumab in combination with TKI in EGFR-positive or ALK-positive NSCLC patients with brain metastases would be particularly valuable.

Young Hak Kim, MD, PhD
Michiaki Mishima, MD, PhD

Department of Respiratory Medicine
Graduate School of Medicine
Kyoto University
Kyoto, Japan

REFERENCES

1. Su YL, Rau KM. Adding bevacizumab to chemotherapy effectively control radioresistant brain metastases in ALK-positive lung adenocarcinoma. *J Thorac Oncol* 2015;10:e21–e22.
2. Socinski MA, Langer CJ, Huang JE, et al. Safety of bevacizumab in patients with non-small-cell lung cancer and brain metastases. *J Clin Oncol* 2009;27:5255–5261.
3. Besse B, Le Moulec S, Mazières J, et al. Bevacizumab in patients with nonsquamous non-small cell lung cancer and asymptomatic, untreated brain metastases (BRAIN): a nonrandomized, phase II study. *Clin Cancer Res* 2015;21:1896–1903.
4. De Braganca KC, Janjigian YY, Azzoli CG, et al. Efficacy and safety of bevacizumab in active brain metastases from non-small cell lung cancer. *J Neurooncol* 2010;100:443–447.
5. Rangachari D, Yamaguchi N, VanderLaan PA, et al. Brain metastases in patients with EGFR-mutated or ALK-rearranged non-small-cell lung cancers. *Lung Cancer* 2015;88:108–111.
6. Noronha V, Joshi A, Gokarn A, et al. The importance of brain metastasis in EGFR mutation positive NSCLC patients. *Chemother Res Pract* 2014;2014:856156.

Address for correspondence: Young Hak Kim, MD, PhD, Department of Respiratory Medicine, Graduate School of Medicine, Kyoto University, 54 Shogoin-Kawaracho, Sakyo-ku, Kyoto 606-8507, Japan. E-mail: ekim@kuhp.kyoto-u.ac.jp

DOI: 10.1097/JTO.0000000000000568

Copyright © 2015 by the International Association for the Study of Lung Cancer

ISSN: 1556-0864/15/1008-0e76




FIGURE 1. Magnetic resonance imaging of the brain (A,B) before and (C,D) after whole brain irradiation, (E,F) after two cycles of chemotherapy of bevacizumab, paclitaxel, and carboplatin.